

Persistent Stiefel-Whitney classes



Raphaël Tinarrage

EPFL, Applied Topology Seminar, 03/11/2020


Persistent homology

2/25 (1/8)


We observe a point cloud X , that we suppose close to a submanifold \mathcal{M} .



Persistent homology in practice:



\mathbb{V}



Filtration $V[X]$



Persistence module $\mathbb{V}[X]$

Barcode $[X]$


Persistent homology

2/25 (2/8)

We observe a point cloud X , that we suppose close to a submanifold \mathcal{M} .




Persistent homology in practice:



Filtration $V[X]$



\mathbb{V}



Persistence module $\mathbb{V}[X]$

Barcode $[X]$

How does $\text{Barcode}[X]$ reveals the homology of \mathcal{M} ?

homotopy type estimation

persistence stability


Persistent homology

Homotopy type estimation


2/25 (3/8)

The Čech filtration of X is the collection $V[X] = (X^t)_{t \geq 0}$ where X^t is the t -thickening of X :


$$X^t = \{y \in \mathbb{R}^n, \exists x \in X, \|x - y\| \leq t\}.$$




$X^0 = X$



$X^{0,1}$



$X^{0,2}$



$X^{0,3}$


Persistent homology

Homotopy type estimation


2/25 (4/8)

The Čech filtration of X is the collection $V[X] = (X^t)_{t \geq 0}$ where X^t is the t -thickening of X :


$$X^t = \{y \in \mathbb{R}^n, \exists x \in X, \|x - y\| \leq t\}.$$




$X^0 = X$





$X^{0,1}$



$X^{0,2}$



$X^{0,3}$



Theorem (Chazal, Cohen-Steiner, Lieutier, 2009)

Let \mathcal{M}, X be subsets of \mathbb{R}^n .

Suppose that $\text{reach}(\mathcal{M}) > 0$ and $d_H(X, \mathcal{M}) \leq \frac{1}{17} \text{reach}(\mathcal{M})$. Let

$$t \in [4d_H(X, \mathcal{M}), \text{reach}(\mathcal{M}) - 3d_H(X, \mathcal{M})].$$

Then X^t and \mathcal{M} are homotopy equivalent.


Persistent homology

Homotopy type estimation


2/25 (5/8)

The Čech filtration of X is the collection $V[X] = (X^t)_{t \geq 0}$ where X^t is the t -thickening of X :


$$X^t = \{y \in \mathbb{R}^n, \exists x \in X, \|x - y\| \leq t\}.$$




$X^0 = X$





$X^{0,1}$



$X^{0,2}$



$X^{0,3}$



Theorem (Niyogi, Smale, Weinberger, 2008)

Let X and \mathcal{M} be subsets of \mathbb{R}^n , with \mathcal{M} a submanifold and $X \subset \mathcal{M}$ finite. Suppose that $\text{reach}(\mathcal{M}) > 0$. Let

$$t \in \left[2d_H(X, \mathcal{M}), \sqrt{\frac{3}{5}} \text{reach}(\mathcal{M}) \right).$$

Then X^t and \mathcal{M} are homotopy equivalent.


Persistent homology

Homotopy type estimation


2/25 (6/8)

The Čech filtration of X is the collection $V[X] = (X^t)_{t \geq 0}$ where X^t is the t -thickening of X :


$$X^t = \{y \in \mathbb{R}^n, \exists x \in X, \|x - y\| \leq t\}.$$




$X^0 = X$



$X^{0,1}$




$X^{0,2}$



$X^{0,3}$

As a consequence, one reads the homology of \mathcal{M} on Barcode[X], on some interval.





Persistent homology

Stability point of view

2/25 (7/8)

Let $V[\mathcal{M}]$ be the Čech filtration of \mathcal{M} .
For every $t \in [0, \text{reach}(\mathcal{M}))$, we have $\mathcal{M}^t \simeq \mathcal{M}$.






Persistent homology

Stability point of view

2/25 (8/8)


Let $V[\mathcal{M}]$ be the Čech filtration of \mathcal{M} .

For every $t \in [0, \text{reach}(\mathcal{M}))$, we have $\mathcal{M}^t \simeq \mathcal{M}$.



Let $\epsilon = d_H(X, \mathcal{M})$.

By stability theorem, $\text{Barcode}[X]$ and $\text{Barcode}[\mathcal{M}]$ are ϵ -close in bottleneck distance.




choose the largest bars!

Other topological invariants


3/25 (1/7)

Persistent homology allows to estimate the *homology* of a space. However, over $\mathbb{Z}/2\mathbb{Z}$, homology may not be fine enough to distinguish between non-homeomorphic spaces.
non-homotopy equivalent spaces.



Torus

$$\mathbb{Z}/2\mathbb{Z}, (\mathbb{Z}/2\mathbb{Z})^2, \mathbb{Z}/2\mathbb{Z}, 0, \dots$$




Klein bottle

$$\mathbb{Z}/2\mathbb{Z}, (\mathbb{Z}/2\mathbb{Z})^2, \mathbb{Z}/2\mathbb{Z}, 0, \dots$$


Other topological invariants

3/25 (2/7)



Persistent homology allows to estimate the *homology* of a space. However, over $\mathbb{Z}/2\mathbb{Z}$, homology may not be fine enough to distinguish between non-homeomorphic spaces.
non-homotopy equivalent spaces.



Torus



Klein bottle


		
	Torus	Klein bottle
$H_i(\mathcal{M}, \mathbb{Z}/2\mathbb{Z}), i \geq 0$	$\mathbb{Z}/2\mathbb{Z}, (\mathbb{Z}/2\mathbb{Z})^2, \mathbb{Z}/2\mathbb{Z}, 0, \dots$	$\mathbb{Z}/2\mathbb{Z}, (\mathbb{Z}/2\mathbb{Z})^2, \mathbb{Z}/2\mathbb{Z}, 0, \dots$
$H_i(\mathcal{M}, \mathbb{Z}/p\mathbb{Z}), i \geq 0$	$\mathbb{Z}/p\mathbb{Z}, (\mathbb{Z}/p\mathbb{Z})^2, \mathbb{Z}/p\mathbb{Z}, 0, \dots$	$\mathbb{Z}/p\mathbb{Z}, \mathbb{Z}/p\mathbb{Z}, 0, 0, \dots$
$H_i(\mathcal{M}, \mathbb{Z}), i \geq 0$	$\mathbb{Z}, \mathbb{Z}^2, \mathbb{Z}, 0, \dots$	$\mathbb{Z}, \mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}, 0, 0, \dots$
$H^*(\mathcal{M}, \mathbb{Z}/2\mathbb{Z})$	$\mathbb{Z}/2\mathbb{Z}[x, y]/\langle x^2, y^2 \rangle$	$\mathbb{Z}/2\mathbb{Z}[x, y]/\langle x^3, y^2, x^2y \rangle$
$w_1(\tau)$	0	x

Other topological invariants


3/25 (3/7)

Persistent homology allows to estimate the *homology* of a space. However, over $\mathbb{Z}/2\mathbb{Z}$, homology may not be fine enough to distinguish between non-homeomorphic spaces.
non-homotopy equivalent spaces.

homology groups
over $\mathbb{Z}/p\mathbb{Z}$



Torus



Klein bottle

$$H_i(\mathcal{M}, \mathbb{Z}/2\mathbb{Z}), i \geq 0$$

$$\mathbb{Z}/2\mathbb{Z}, (\mathbb{Z}/2\mathbb{Z})^2, \mathbb{Z}/2\mathbb{Z}, 0, \dots$$

$$\mathbb{Z}/2\mathbb{Z}, (\mathbb{Z}/2\mathbb{Z})^2, \mathbb{Z}/2\mathbb{Z}, 0, \dots$$

$$H_i(\mathcal{M}, \mathbb{Z}/p\mathbb{Z}), i \geq 0$$

$$\mathbb{Z}/p\mathbb{Z}, (\mathbb{Z}/p\mathbb{Z})^2, \mathbb{Z}/p\mathbb{Z}, 0, \dots$$

$$\mathbb{Z}/p\mathbb{Z}, \mathbb{Z}/p\mathbb{Z}, 0, 0, \dots$$

$$H_i(\mathcal{M}, \mathbb{Z}), i \geq 0$$

$$\mathbb{Z}, \mathbb{Z}^2, \mathbb{Z}, 0, \dots$$

$$\mathbb{Z}, \mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}, 0, 0, \dots$$

$$H^*(\mathcal{M}, \mathbb{Z}/2\mathbb{Z})$$

$$\mathbb{Z}/2\mathbb{Z}[x, y] / \langle x^2, y^2 \rangle$$

$$\mathbb{Z}/2\mathbb{Z}[x, y] / \langle x^3, y^2, x^2y \rangle$$

$$w_1(\tau)$$

$$0$$


$$x$$

Other topological invariants


3/25 (4/7)

Persistent homology allows to estimate the *homology* of a space. However, over $\mathbb{Z}/2\mathbb{Z}$, homology may not be fine enough to distinguish between non-homeomorphic spaces.
non-homotopy equivalent spaces.

homology groups
over \mathbb{Z}



Torus



Klein bottle


$H_i(\mathcal{M}, \mathbb{Z}/2\mathbb{Z}), i \geq 0$	$\mathbb{Z}/2\mathbb{Z}, (\mathbb{Z}/2\mathbb{Z})^2, \mathbb{Z}/2\mathbb{Z}, 0, \dots$	$\mathbb{Z}/2\mathbb{Z}, (\mathbb{Z}/2\mathbb{Z})^2, \mathbb{Z}/2\mathbb{Z}, 0, \dots$
$H_i(\mathcal{M}, \mathbb{Z}/p\mathbb{Z}), i \geq 0$	$\mathbb{Z}/p\mathbb{Z}, (\mathbb{Z}/p\mathbb{Z})^2, \mathbb{Z}/p\mathbb{Z}, 0, \dots$	$\mathbb{Z}/p\mathbb{Z}, \mathbb{Z}/p\mathbb{Z}, 0, 0, \dots$
$H_i(\mathcal{M}, \mathbb{Z}), i \geq 0$	$\mathbb{Z}, \mathbb{Z}^2, \mathbb{Z}, 0, \dots$	$\mathbb{Z}, \mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}, 0, 0, \dots$
$H^*(\mathcal{M}, \mathbb{Z}/2\mathbb{Z})$	$\mathbb{Z}/2\mathbb{Z}[x, y]/\langle x^2, y^2 \rangle$	$\mathbb{Z}/2\mathbb{Z}[x, y]/\langle x^3, y^2, x^2y \rangle$
$w_1(\tau)$	0	x

Other topological invariants


3/25 (5/7)

Persistent homology allows to estimate the *homology* of a space. However, over $\mathbb{Z}/2\mathbb{Z}$, homology may not be fine enough to distinguish between non-homeomorphic spaces.
non-homotopy equivalent spaces.

cohomology
algebra over $\mathbb{Z}/2\mathbb{Z}$



Torus



Klein bottle

$$H_i(\mathcal{M}, \mathbb{Z}/2\mathbb{Z}), i \geq 0$$

$$\mathbb{Z}/2\mathbb{Z}, (\mathbb{Z}/2\mathbb{Z})^2, \mathbb{Z}/2\mathbb{Z}, 0, \dots$$

$$\mathbb{Z}/2\mathbb{Z}, (\mathbb{Z}/2\mathbb{Z})^2, \mathbb{Z}/2\mathbb{Z}, 0, \dots$$

$$H_i(\mathcal{M}, \mathbb{Z}/p\mathbb{Z}), i \geq 0$$

$$\mathbb{Z}/p\mathbb{Z}, (\mathbb{Z}/p\mathbb{Z})^2, \mathbb{Z}/p\mathbb{Z}, 0, \dots$$

$$\mathbb{Z}/p\mathbb{Z}, \mathbb{Z}/p\mathbb{Z}, 0, 0, \dots$$

$$H_i(\mathcal{M}, \mathbb{Z}), i \geq 0$$

$$\mathbb{Z}, \mathbb{Z}^2, \mathbb{Z}, 0, \dots$$

$$\mathbb{Z}, \mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}, 0, 0, \dots$$

$$H^*(\mathcal{M}, \mathbb{Z}/2\mathbb{Z})$$

$$\mathbb{Z}/2\mathbb{Z}[x, y] / \langle x^2, y^2 \rangle$$

$$\mathbb{Z}/2\mathbb{Z}[x, y] / \langle x^3, y^2, x^2 y \rangle$$

$$w_1(\tau)$$

$$0$$


$$x$$

Other topological invariants


3/25 (6/7)

Persistent homology allows to estimate the *homology* of a space. However, over $\mathbb{Z}/2\mathbb{Z}$, homology may not be fine enough to distinguish between non-homeomorphic spaces.
non-homotopy equivalent spaces.

first Stiefel-Whitney
class of tangent bundle



Torus



Klein bottle

$$H_i(\mathcal{M}, \mathbb{Z}/2\mathbb{Z}), i \geq 0$$

$$\mathbb{Z}/2\mathbb{Z}, (\mathbb{Z}/2\mathbb{Z})^2, \mathbb{Z}/2\mathbb{Z}, 0, \dots$$

$$\mathbb{Z}/2\mathbb{Z}, (\mathbb{Z}/2\mathbb{Z})^2, \mathbb{Z}/2\mathbb{Z}, 0, \dots$$

$$H_i(\mathcal{M}, \mathbb{Z}/p\mathbb{Z}), i \geq 0$$

$$\mathbb{Z}/p\mathbb{Z}, (\mathbb{Z}/p\mathbb{Z})^2, \mathbb{Z}/p\mathbb{Z}, 0, \dots$$

$$\mathbb{Z}/p\mathbb{Z}, \mathbb{Z}/p\mathbb{Z}, 0, 0, \dots$$

$$H_i(\mathcal{M}, \mathbb{Z}), i \geq 0$$

$$\mathbb{Z}, \mathbb{Z}^2, \mathbb{Z}, 0, \dots$$

$$\mathbb{Z}, \mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}, 0, 0, \dots$$

$$H^*(\mathcal{M}, \mathbb{Z}/2\mathbb{Z})$$

$$\mathbb{Z}/2\mathbb{Z}[x, y] / \langle x^2, y^2 \rangle$$

$$\mathbb{Z}/2\mathbb{Z}[x, y] / \langle x^3, y^2, x^2 y \rangle$$

$$w_1(\tau)$$

$$0$$


$$x$$

Other topological invariants


3/25 (7/7)

Persistent homology allows to estimate the *homology* of a space. However, over $\mathbb{Z}/2\mathbb{Z}$, homology may not be fine enough to distinguish between non-homeomorphic spaces.
non-homotopy equivalent spaces.

first Stiefel-Whitney
class of tangent bundle



Torus



Klein bottle

Aim of this talk:

Building a persistent framework for Stiefel-Whitney classes, with consistency and stability inspired from persistent homology.

$w_1(\tau)$

0

x

I - Stiefel-Whitney classes

II - Persistent Stiefel-Whitney classes

III - Algorithmic considerations



Definition:

A vector bundle (of dimension d) over X is a surjection $\pi : E \rightarrow X$, with E a topological space, such that:

- the fibers $\pi^{-1}(\{x\})$, $x \in X$, are vector spaces of dimension d ,
- π satisfies a local triviality condition.

Local triviality condition: for all $x \in X$, there exists a neighborhood $U \subset X$ and a homeomorphism $h: U \times \mathbb{R}^d \rightarrow \pi^{-1}(U)$ such that for all $y \in U$, $h(y, \cdot)$ is an isomorphism of vector spaces.

Normal bundle
of the circle



Möbius strip
(universal
bundle)



Definition:

A vector bundle (of dimension d) over X is a surjection $\pi : E \rightarrow X$, with E a topological space, such that:

- the fibers $\pi^{-1}(\{x\})$, $x \in X$, are vector spaces of dimension d ,
- π satisfies a local triviality condition.

Local triviality condition: for all $x \in X$, there exists a neighborhood $U \subset X$ and a homeomorphism $h: U \times \mathbb{R}^d \rightarrow \pi^{-1}(U)$ such that for all $y \in U$, $h(y, \cdot)$ is an isomorphism of vector spaces.

Tangent bundle
of sphere



Normal bundle
of the torus

Stiefel-Whitney classes (axioms)

6/25 (1/2)

For every vector bundle $\pi: E \rightarrow X$, there exists a sequence of cohomology classes

$$w_0(\pi) \in H^0(X, \mathbb{Z}/2\mathbb{Z}),$$

$$w_1(\pi) \in H^1(X, \mathbb{Z}/2\mathbb{Z}),$$

$$w_2(\pi) \in H^2(X, \mathbb{Z}/2\mathbb{Z}),$$

$$w_3(\pi) \in H^3(X, \mathbb{Z}/2\mathbb{Z}),$$

...

that satisfy the following axioms:

- **Axiom 1:** $w_0(\pi)$ is equal to $1 \in H^0(X, \mathbb{Z}/2\mathbb{Z})$, and if π is of dimension d then $w_i(\pi) = 0$ for $i > d$.
- **Axiom 2:** if $f: \pi \rightarrow \rho$ is a bundle map, then $w_i(\pi) = f^*(w_i(\rho))$, where $f^*: H^*(X) \leftarrow H^*(Y)$ is the map induced in cohomology by f .
- **Axiom 3:** if π, ρ are vector bundles over the same base space X , then for all $k \in \mathbb{N}$, $w_k(\pi \oplus \rho) = \sum_{i=0}^k w_i(\pi) \smile w_{k-i}(\rho)$ (cup product).
- **Axiom 4:** $w_1(\gamma_1^1) \neq 0$, where γ_1^1 denotes the Möbius strip bundle over the circle.

For every vector bundle $\pi: E \rightarrow X$, there exists a sequence of cohomology classes

$$w_0(\pi) \in H^0(X, \mathbb{Z}/2\mathbb{Z}),$$

$$w_1(\pi) \in H^1(X, \mathbb{Z}/2\mathbb{Z}),$$

$$w_2(\pi) \in H^2(X, \mathbb{Z}/2\mathbb{Z}),$$

$$w_3(\pi) \in H^3(X, \mathbb{Z}/2\mathbb{Z}),$$

...

Basic properties:

- If two bundles are isomorphic, then their Stiefel-Whitney classes are equal.
- If π admits a (nowhere vanishing) section, then $w_d(\pi) = 0$.
- If π admits k independent (nowhere vanishing) sections, then $w_d(\pi) = \dots w_{d-k+1}(\pi) = 0$.

Topological information:


- If τ is the tangent bundle of a manifold \mathcal{M} , then \mathcal{M} is orientable if and only if $w_1(\tau) = 0$.
- \mathcal{M} admits a spin structure if and only if $w_1(\tau) = 0$ and $w_2(\tau) = 0$.

The Grassmann manifolds


7/25 (1/3)

Let $d, n \geq 1$.

The *Grassmannian* $\mathcal{G}_d(\mathbb{R}^n)$ is the set of d -dimensional linear subspaces of \mathbb{R}^n . It can be endowed with a manifold structure, of dimension $d(n - d)$.





$\mathcal{G}_1(\mathbb{R}^2)$



$\mathcal{G}_1(\mathbb{R}^3) \simeq \mathcal{G}_2(\mathbb{R}^3)$

A construction of $\mathcal{G}_1(\mathbb{R}^2)$:




The Grassmann manifolds

7/25 (2/3)


Let $d, n \geq 1$.

The *Grassmannian* $\mathcal{G}_d(\mathbb{R}^n)$ is the set of d -dimensional linear subspaces of \mathbb{R}^n .

It can be endowed with a manifold structure, of dimension $d(n - d)$.



$\mathcal{G}_1(\mathbb{R}^2)$



$\mathcal{G}_1(\mathbb{R}^3) \simeq \mathcal{G}_2(\mathbb{R}^3)$

Let \mathbb{R}^∞ denotes the space of sequences of real numbers that are zero from some point. We can also define the *infinite Grassmannian* $\mathcal{G}_d(\mathbb{R}^\infty)$.

The infinite Grassmannian has $\mathbb{Z}/2\mathbb{Z}$ -cohomology

$$H^*(\mathcal{G}_d(\mathbb{R}^\infty)) = \mathbb{Z}/2\mathbb{Z}[w_1, \dots, w_d]$$

where w_i has degree i .

In particular, $H^*(\mathcal{G}_1(\mathbb{R}^\infty)) = \mathbb{Z}/2\mathbb{Z}[w_1]$.


The Grassmann manifolds

7/25 (3/3)


Let $d, n \geq 1$.

The *Grassmannian* $\mathcal{G}_d(\mathbb{R}^n)$ is the set of d -dimensional linear subspaces of \mathbb{R}^n .

It can be endowed with a manifold structure, of dimension $d(n - d)$.



$\mathcal{G}_1(\mathbb{R}^2)$



$\mathcal{G}_1(\mathbb{R}^3) \simeq \mathcal{G}_2(\mathbb{R}^3)$

Let $M(\mathbb{R}^n)$ be the space of $n \times n$ matrices.

For every linear subspace $T \subset \mathbb{R}^n$, let p_T denotes the orthogonal projection matrix on T .

The application $T \in \mathcal{G}_d(\mathbb{R}^n) \longmapsto p_T \in M(\mathbb{R}^n)$ is an embedding.

Hence $\mathcal{G}_d(\mathbb{R}^n)$ can be seen as a submanifold of $M(\mathbb{R}^n)$.

Vector bundles (2nd definition)


8/25

Correspondence vector bundles / classifying maps:

Let X is a topological space. From any continuous map $\xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^n)$, we can build a d -dimensional vector bundle structure on X .

Conversely, for any vector bundle $\pi: E \rightarrow X$, there exists a corresponding map $\xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^\infty)$, called a *classifying map*.

Moreover, if X is compact, we can choose $\xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^m)$ for m large enough.



(Second) Definition:

A vector bundle over X is a continuous map $\xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^\infty)$ or $\xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^m)$.

Stiefel-Whitney classes (construction) 9/25 (1/2)

Let $\xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^\infty)$ be a vector bundle, and
 $\xi^*: H^*(X, \mathbb{Z}/2\mathbb{Z}) \leftarrow H^*(\mathcal{G}_d(\mathbb{R}^\infty), \mathbb{Z}/2\mathbb{Z})$ the map induced in cohomology.

Recall that $H^*(\mathcal{G}_d(\mathbb{R}^\infty)) = \mathbb{Z}/2\mathbb{Z}[w_1, \dots, w_d]$.


The Stiefel-Whitney classes of the vector bundle $\xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^\infty)$ can be defined as

$$w_0(\xi) = \xi^*(\omega_0)$$

$$w_1(\xi) = \xi^*(\omega_1)$$

$$w_2(\xi) = \xi^*(\omega_2)$$

...



Stiefel-Whitney classes (construction) 9/25 (2/2)

Let $\xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^\infty)$ be a vector bundle, and
 $\xi^*: H^*(X, \mathbb{Z}/2\mathbb{Z}) \leftarrow H^*(\mathcal{G}_d(\mathbb{R}^\infty), \mathbb{Z}/2\mathbb{Z})$ the map induced in cohomology.

Recall that $H^*(\mathcal{G}_d(\mathbb{R}^\infty)) = \mathbb{Z}/2\mathbb{Z}[w_1, \dots, w_d]$.


The Stiefel-Whitney classes of the vector bundle $\xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^\infty)$ can be defined as

$$w_0(\xi) = \xi^*(\omega_0)$$

$$w_1(\xi) = \xi^*(\omega_1)$$

$$w_2(\xi) = \xi^*(\omega_2)$$

...



I - Stiefel-Whitney classes

II - Persistent Stiefel-Whitney classes

III - Algorithmic considerations


Adopting a persistent viewpoint

11/25 (1/3)

Sampling model for vector bundles:

Let $n, m, d > 0$.

We observe $\left\{ \begin{array}{l} \text{a point cloud } X \subset \mathbb{R}^n \\ \text{and a map } \xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^m). \end{array} \right.$




Adopting a persistent viewpoint

11/25 (2/3)

Sampling model for vector bundles:

Let $n, m, d > 0$.


We observe $\left\{ \begin{array}{l} \text{a point cloud } X \subset \mathbb{R}^n \\ \text{and a map } \xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^m). \end{array} \right.$



Defining a vector bundle filtration:

Let $(X^t)_{t \geq 0}$ be the Čech filtration of X .

We want to define maps $\xi^t: X^t \rightarrow \mathcal{G}_d(\mathbb{R}^m)$.




Adopting a persistent viewpoint

11/25 (3/3)

Sampling model for vector bundles:

Let $n, m, d > 0$.


We observe $\left\{ \begin{array}{l} \text{a point cloud } X \subset \mathbb{R}^n \\ \text{and a map } \xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^m). \end{array} \right.$



Defining a vector bundle filtration:

Let $(X^t)_{t \geq 0}$ be the Čech filtration of X .

We want to define maps $\xi^t: X^t \rightarrow \mathcal{G}_d(\mathbb{R}^m)$.



Nothing interesting to do here...

A persistent viewpoint (2nd attempt) 12/25 (1/3)

Sampling model for vector bundles:

Let $n, m, d > 0$.

We observe $\left| \begin{array}{l} \text{a point cloud } X \subset \mathbb{R}^n \\ \text{and a map } \xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^m). \end{array} \right. \iff \text{a point cloud } \check{X} \subset \mathbb{R}^n \times \mathcal{G}_d(\mathbb{R}^m).$

$$\check{X} = \{(x, \xi(x)), x \in X\}$$

A persistent viewpoint (2nd attempt) 12/25 (2/3)

Sampling model for vector bundles:

Let $n, m, d > 0$.

We observe $\left| \begin{array}{l} \text{a point cloud } X \subset \mathbb{R}^n \\ \text{and a map } \xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^m). \end{array} \right. \iff \text{a point cloud } \check{X} \subset \mathbb{R}^n \times \mathcal{G}_d(\mathbb{R}^m).$

$$\check{X} = \{(x, \xi(x)), x \in X\}$$

- By embedding $\mathcal{G}_d(\mathbb{R}^m) \hookrightarrow M(\mathbb{R}^m)$, we can see \check{X} as a subset of $\mathbb{R}^n \times M(\mathbb{R}^m)$.
- Let $(\check{X}^t)_{t \geq 0}$ be the Čech filtration of \check{X} in the ambient space $\mathbb{R}^n \times M(\mathbb{R}^m)$, endowed with the metric $\|(x, A)\| = \sqrt{\|x\|_2^2 + \|A\|_F^2}$.
- We can define extended maps ξ^t as follows:

$$\begin{aligned} \xi^t: \quad \check{X}^t &\longrightarrow \mathcal{G}_d(\mathbb{R}^m) \\ (x, A) &\longmapsto \text{proj}(A, \mathcal{G}_d(\mathbb{R}^m)) \end{aligned}$$

A persistent viewpoint (2nd attempt) 12/25 (3/3)

Sampling model for vector bundles:

Let $n, m, d > 0$.

We observe $\left| \begin{array}{l} \text{a point cloud } X \subset \mathbb{R}^n \\ \text{and a map } \xi: X \rightarrow \mathcal{G}_d(\mathbb{R}^m). \end{array} \right. \iff \text{a point cloud } \check{X} \subset \mathbb{R}^n \times \mathcal{G}_d(\mathbb{R}^m).$

$$\check{X} = \{(x, \xi(x)), x \in X\}$$

- By embedding $\mathcal{G}_d(\mathbb{R}^m) \hookrightarrow M(\mathbb{R}^m)$, we can see \check{X} as a subset of $\mathbb{R}^n \times M(\mathbb{R}^m)$.
- Let $(\check{X}^t)_{t \geq 0}$ be the Čech filtration of \check{X} in the ambient space $\mathbb{R}^n \times M(\mathbb{R}^m)$, endowed with the metric $\|(x, A)\| = \sqrt{\|x\|_2^2 + \|A\|_F^2}$.
- We can define extended maps ξ^t as follows:

$$\begin{aligned} \xi^t: \quad \check{X}^t &\longrightarrow \mathcal{G}_d(\mathbb{R}^m) \\ (x, A) &\longmapsto \text{proj}(A, \mathcal{G}_d(\mathbb{R}^m)) \end{aligned}$$

Definition:

The data of $(\check{X}^t)_{t \geq 0}$ and $(\xi^t: \check{X}^t \rightarrow \mathcal{G}_d(\mathbb{R}^m))_{t \geq 0}$ is called the *Čech bundle filtration* of \check{X} .

Let $\left| \begin{array}{l} \check{X} \subset \mathbb{R}^n \times \mathcal{G}_d(\mathbb{R}^m), \\ (\check{X}^t)_t, (\xi^t)_t \text{ its Čech bundle filtration,} \\ i \geq 0. \end{array} \right.$

For every $t \geq 0$, we have the i^{th} Stiefel-Whitney class of (\check{X}^t, ξ^t) :

$$w_i(\xi^t) = (\xi^t)^*(w_i),$$

where $(\xi^t)^*: H^*(\check{X}^t) \leftarrow H^*(\mathcal{G}_d(\mathbb{R}^m))$.

Definition:

The i^{th} persistent Stiefel-Whitney class of \check{X} is the collection $(w_i(\xi^t))_{t \geq 0}$.

Let $\left| \begin{array}{l} \check{X} \subset \mathbb{R}^n \times \mathcal{G}_d(\mathbb{R}^m), \\ (\check{X}^t)_t, (\xi^t)_t \text{ its Čech bundle filtration,} \\ i \geq 0. \end{array} \right.$

For every $t \geq 0$, we have the i^{th} Stiefel-Whitney class of (\check{X}^t, ξ^t) :

$$w_i(\xi^t) = (\xi^t)^*(w_i),$$

where $(\xi^t)^*: H^*(\check{X}^t) \leftarrow H^*(\mathcal{G}_d(\mathbb{R}^m))$.

Definition:

The i^{th} persistent Stiefel-Whitney class of \check{X} is the collection $(w_i(\xi^t))_{t \geq 0}$.

Issue: ξ^t is not well-defined for every $t \geq 0 \dots$

The extended maps ξ^t are defined as

$$\begin{aligned}\xi^t: \quad \check{X}^t &\longrightarrow \mathcal{G}_d(\mathbb{R}^m) \\ (x, A) &\longmapsto \text{proj}(A, \mathcal{G}_d(\mathbb{R}^m))\end{aligned}$$

But $\text{proj}(A, \mathcal{G}_d(\mathbb{R}^m))$ does not make sense if A lies in the medial axis of $\mathcal{G}_d(\mathbb{R}^m)$.

→ There exists a maximal value t^{\max} such that for all $t \in [0, t^{\max})$, the maps ξ^t are well-defined.

The extended maps ξ^t are defined as

$$\begin{aligned}\xi^t: \quad \check{X}^t &\longrightarrow \mathcal{G}_d(\mathbb{R}^m) \\ (x, A) &\longmapsto \text{proj}(A, \mathcal{G}_d(\mathbb{R}^m))\end{aligned}$$

But $\text{proj}(A, \mathcal{G}_d(\mathbb{R}^m))$ does not make sense if A lies in the medial axis of $\mathcal{G}_d(\mathbb{R}^m)$.

→ There exists a maximal value t^{\max} such that for all $t \in [0, t^{\max})$, the maps ξ^t are well-defined.

Lemma

For any $A \in M(\mathbb{R}^m)$, let A^s denote the matrix $A^s = \frac{1}{2}(A + {}^tA)$, and let $\lambda_1(A^s), \dots, \lambda_n(A^s)$ be the eigenvalues of A^s in decreasing order.

The distance from A to $\text{med}(\mathcal{G}_d(\mathbb{R}^m))$ is $\frac{\sqrt{2}}{2} |\lambda_d(A^s) - \lambda_{d+1}(A^s)|$.

The persistent Stiefel-Whitney class $(w_i(\xi^t))_t$ is defined for every $t \in [0, t^{\max})$.

Let $\check{X} \subset \mathbb{R}^n \times M(\mathbb{R}^m)$, and $w_i(\check{X})$ its i^{th} persistent Stiefel-Whitney class.

Definition

The *lifebar* of the persistent Stiefel-Whitney class $w_i(\check{X})$ is the set

$$\{t \in [0, t^{\max}), w_i(\xi^t) \neq 0\}.$$



the lifebar is an interval!

Let $\check{X} \subset \mathbb{R}^n \times M(\mathbb{R}^m)$, and $w_i(\check{X})$ its i^{th} persistent Stiefel-Whitney class.

Definition




The *lifebar* of the persistent Stiefel-Whitney class $w_i(\check{X})$ is the set

$$\{t \in [0, t^{\max}), w_i(\xi^t) \neq 0\}.$$









the lifebar is an interval!


Example: lifebars of first persistent Stiefel-Whitney classes



Theorem

If two subsets $\check{X}, \check{Y} \subset \mathbb{R}^n \times M(\mathbb{R}^m)$ satisfies $d_H(\check{X}, \check{Y}) \leq \epsilon$, then for all $i \geq 0$, the lifebars of their i^{th} Stiefel-Whitney classes are ϵ -close.


 $w_1(\check{X})$

 $w_1(\check{Y})$

 $w_1(\check{X})$

 $w_1(\check{Y})$


Consistency

17/25 (1/2)



If $u: \mathcal{M}_0 \rightarrow \mathcal{M} \subset \mathbb{R}^n$ is an immersion and $\xi: \mathcal{M}_0 \rightarrow \mathcal{G}_d(\mathbb{R}^m)$ a vector bundle, consider the set

$$\check{\mathcal{M}} = \{(u(x_0), \xi(x_0)), x_0 \in \mathcal{M}_0\} \subset \mathbb{R}^n \times \mathbf{M}(\mathbb{R}^m).$$


Theorem

Let $X \subset \mathbb{R}^n \times \mathbf{M}(\mathbb{R}^m)$ be any subset such that $d_H(X, \check{\mathcal{M}}) \leq \epsilon$. Then for every $t \in [4\epsilon, \text{reach}(\check{\mathcal{M}}) - 3\epsilon]$, the composition of inclusions $\mathcal{M}_0 \hookrightarrow \check{\mathcal{M}} \hookrightarrow X^t$ induces an isomorphism $H^*(\mathcal{M}_0) \leftarrow H^*(X^t)$ which sends the i^{th} persistent Stiefel-Whitney class $w_i^t(X)$ of the Čech bundle filtration of X to the i^{th} Stiefel-Whitney class of (\mathcal{M}_0, p) .

Normal bundle of the torus



Normal bundle of the Klein bottle



Consistency

17/25 (2/2)


If $u: \mathcal{M}_0 \rightarrow \mathcal{M} \subset \mathbb{R}^n$ is an immersion and $\xi: \mathcal{M}_0 \rightarrow \mathcal{G}_d(\mathbb{R}^m)$ a vector bundle, consider the set

$$\check{\mathcal{M}} = \{(u(x_0), \xi(x_0)), x_0 \in \mathcal{M}_0\} \subset \mathbb{R}^n \times \mathbf{M}(\mathbb{R}^m).$$


Theorem

Let $X \subset \mathbb{R}^n \times \mathbf{M}(\mathbb{R}^m)$ be any subset such that $d_H(X, \check{\mathcal{M}}) \leq \epsilon$. Then for every $t \in [4\epsilon, \text{reach}(\check{\mathcal{M}}) - 3\epsilon]$, the composition of inclusions $\mathcal{M}_0 \hookrightarrow \check{\mathcal{M}} \hookrightarrow X^t$ induces an isomorphism $H^*(\mathcal{M}_0) \leftarrow H^*(X^t)$ which sends the i^{th} persistent Stiefel-Whitney class $w_i^t(X)$ of the Čech bundle filtration of X to the i^{th} Stiefel-Whitney class of (\mathcal{M}_0, p) .

Normal bundle of the torus




Normal bundle of the Klein bottle



orientable

non-orientable



I - Stiefel-Whitney classes

II - Persistent Stiefel-Whitney classes

III - Algorithmic considerations

Let $\check{X} \subset \mathbb{R}^n \times \mathcal{G}_d(\mathbb{R}^m)$ or $\check{X} \subset \mathbb{R}^n \times M(\mathbb{R}^m)$,
 $(\check{X}^t)_t, (\xi^t)_t$ its Čech bundle filtration,
 $(w_i(\xi^t))_t$ its i^{th} persistent Stiefel-Whitney class.

$$\begin{array}{ccc} \xi^t: & \check{X}^t & \longrightarrow \mathcal{G}_d(\mathbb{R}^m) \\ (\xi^t)^*: & H^*(\check{X}^t) & \longleftarrow H^*(\mathcal{G}_d(\mathbb{R}^m)) \\ & w_i(\xi^t) & \longleftarrow w_i \end{array}$$

Problem:

Compute $w_i(\xi^t)$ on a computer.

Let $\check{X} \subset \mathbb{R}^n \times \mathcal{G}_d(\mathbb{R}^m)$ or $\check{X} \subset \mathbb{R}^n \times M(\mathbb{R}^m)$,
 $(\check{X}^t)_t, (\xi^t)_t$ its Čech bundle filtration,
 $(w_i(\xi^t))_t$ its i^{th} persistent Stiefel-Whitney class.


$$\begin{array}{ccc} \xi^t: & \check{X}^t & \longrightarrow \mathcal{G}_d(\mathbb{R}^m) \\ (\xi^t)^*: & H^*(\check{X}^t) & \longleftarrow H^*(\mathcal{G}_d(\mathbb{R}^m)) \\ & w_i(\xi^t) & \longleftarrow w_i \end{array}$$

Problem:

Compute $w_i(\xi^t)$ on a computer.

Suppose that we have triangulations S^t of \check{X}^t and G of $\mathcal{G}_d(\mathbb{R}^m)$.

nerve of the union of balls 

 see later

Let $\check{X} \subset \mathbb{R}^n \times \mathcal{G}_d(\mathbb{R}^m)$ or $\check{X} \subset \mathbb{R}^n \times M(\mathbb{R}^m)$,
 $(\check{X}^t)_t, (\xi^t)_t$ its Čech bundle filtration,
 $(w_i(\xi^t))_t$ its i^{th} persistent Stiefel-Whitney class.

$$\begin{array}{ccc} \xi^t: & \check{X}^t & \longrightarrow \mathcal{G}_d(\mathbb{R}^m) \\ (\xi^t)^*: & H^*(\check{X}^t) & \longleftarrow H^*(\mathcal{G}_d(\mathbb{R}^m)) \\ & w_i(\xi^t) & \longleftarrow w_i \end{array}$$

Problem:

Compute $w_i(\xi^t)$ on a computer.

Suppose that we have triangulations S^t of \check{X}^t and G of $\mathcal{G}_d(\mathbb{R}^m)$.

Denote their topological realizations $|S^t|$ and $|G|$.


$$\begin{array}{ccccc} \check{X}^t & \xrightarrow{\xi^t} & \mathcal{G}_d(\mathbb{R}^m) & & \\ \uparrow \wr & & \uparrow \wr & & \\ |S^t| & \xrightarrow{\quad\quad\quad} & |G| & \xrightarrow{\quad\quad\quad} & G \end{array}$$

$S^t \xrightarrow{\quad\quad\quad ? \quad\quad\quad} G$

We look for a simplicial map $p^t: S^t \rightarrow G$ that 'corresponds to' ξ^t .


Star condition


20/25 (1/3)



The map g satisfies the *star condition* if:

for every vertex $v \in S^t$, there exists a vertex $w \in G$ such that $g(|\overline{\text{St}}(v)|) \subseteq |\text{St}(w)|$.





The map g satisfies the *star condition* if:


for every vertex $v \in S^t$, there exists a vertex $w \in G$ such that $g(|\overline{\text{St}}(v)|) \subseteq |\text{St}(w)|$.


If this is the case, let $f: S^t \rightarrow G$ be any map between vertex sets such that:

for every vertex $v \in S^t$, we have $g(|\overline{\text{St}}(v)|) \subseteq |\text{St}(f(v))|$.

Such a map f is called a *simplicial approximation* to g . It is a simplicial map.

Its topological realization $|f|$ is homotopic to g .





The map g satisfies the *star condition* if:

for every vertex $v \in S^t$, there exists a vertex $w \in G$ such that $g(|\overline{\text{St}}(v)|) \subseteq |\text{St}(w)|$.

If this is the case, let $f: S^t \rightarrow G$ be any map between vertex sets such that:


for every vertex $v \in S^t$, we have $g(|\overline{\text{St}}(v)|) \subseteq |\text{St}(f(v))|$.

Such a map f is called a *simplicial approximation* to g . It is a simplicial map.

Its topological realization $|f|$ is homotopic to g .

Remark:


If g does not satisfy the star condition, we can apply barycentric subdivisions to S^t .



Weak star condition

21/25 (1/2)


In practice, we cannot check whether g satisfies the star condition...




$$g(|\overline{\text{St}}(v)|) \subseteq |\text{St}(w)|?$$

The map g satisfies the *weak star condition* if:

for every vertex $v \in S^t$, there exists a vertex $w \in G$ such that
 $g(|\text{vertices}(\overline{\text{St}}(v))|) \subseteq |\text{St}(w)|$.



In practice, we cannot check whether g satisfies the star condition...



$$g(|\overline{\text{St}}(\textcolor{red}{v})|) \subseteq |\text{St}(\textcolor{violet}{w})|?$$

The map g satisfies the *weak star condition* if:

for every vertex $\textcolor{red}{v} \in S^t$, there exists a vertex $\textcolor{violet}{w} \in G$ such that $g(|\text{vertices}(\overline{\text{St}}(\textcolor{red}{v}))|) \subseteq |\text{St}(\textcolor{violet}{w})|$.

If this is the case, let $f: S^t \rightarrow G$ be any map between vertex sets such that:

for every vertex $\textcolor{red}{v} \in S^t$, we have $g(|\text{vertices}(\overline{\text{St}}(\textcolor{red}{v}))|) \subseteq |\text{St}(\textcolor{violet}{f}(\textcolor{red}{v}))|$.

Such a map f is called a *weak simplicial approximation* to g . It is a simplicial map.

Proposition:


If S^t is subdivided enough, then any weak simplicial approximation is a simplicial approximation.

Triangulations of the Grassmannian 22/25 (1/2)

The Grassmannian $\mathcal{G}_d(\mathbb{R}^n)$ has a well-known CW-complex structure.

However, I had some troubles finding explicit triangulations of $\mathcal{G}_d(\mathbb{R}^n)$.

What is known: triangulations of $\mathcal{G}_1(\mathbb{R}^n)$, the *projective spaces*.




Triangulations of the Grassmannian 22/25 (2/2)

The Grassmannian $\mathcal{G}_d(\mathbb{R}^n)$ has a well-known CW-complex structure.

However, I had some troubles finding explicit triangulations of $\mathcal{G}_d(\mathbb{R}^n)$.

What is known: triangulations of $\mathcal{G}_1(\mathbb{R}^n)$, the *projective spaces*.



In practice, we will only consider the case $d = 1$.

An algorithm for $w_1(\xi^t)$, t fixed

23/25

Consider the map $\xi^t: \check{X}^t \rightarrow \mathcal{G}_1(\mathbb{R}^m)$. We want to compute $w_1(\xi^t) = (\xi^t)^*(w_1)$.

Reminder: $H^1(\mathcal{G}_1(\mathbb{R}^m)) = \langle w_1 \rangle \simeq \mathbb{Z}/2\mathbb{Z}$.

We have to find the image of $(\xi^t)^*: H^1(X^t) \leftarrow H^1(\mathcal{G}_1(\mathbb{R}^m))$

- Compute a triangulation S^t of \check{X}^t
- Compute a triangulation G of $\mathcal{G}_1(\mathbb{R}^m)$
- Check whether ξ^t satisfies the weak star condition
- If not, subdivide barycentric
- Compute a weak simplicial approximation f to ξ^t
- Compute the induced map in simplicial cohomology $f^*: H^1(S^t) \leftarrow H^1(G)$

—————→ The image of f^* is $w_1(\xi^t)$
(seen in simplicial cohomology)

Let $\left| \begin{array}{l} \check{X} \subset \mathbb{R}^n \times \mathcal{G}_d(\mathbb{R}^m), \\ (\check{X}^t)_t, (\xi^t)_t \text{ its Čech bundle filtration,} \\ (w_i(\xi^t))_t \text{ its } i^{\text{th}} \text{ persistent Stiefel-Whitney class.} \end{array} \right.$

We have seen how to compute $w_1(\xi^t)$, t fixed.

Recall that the lifebar of $w_1(X)$ is the set

$$\{t < t_{\max}, w_1(\xi^t) \neq 0\}.$$




Three possibilities for computing the lifebar:

- Compute $w_1(\xi^t)$ for several values of t , and check whether $w_1(\xi^t) = 0$ (dichotomic search)
- Use the persistent image algorithm of [Cohen-Steiner, Edelsbrunner, Harer, Morozov]
- Use the formula on the next page

Reminder: $H^1(\mathcal{G}_1(\mathbb{R}^m)) = \mathbb{Z}/2\mathbb{Z}$.

We have to find the image of $(\xi^t)^*: H^1(\check{X}^t) \leftarrow H^1(\mathcal{G}_1(\mathbb{R}^m))$

Let $C(\xi^t)$ be **the mapping cone** of $\xi^t: \check{X}^t \rightarrow \mathcal{G}_1(\mathbb{R}^m)$.



We have a long exact sequence

$$\dots \longrightarrow H^k(\check{X}^t) \longrightarrow H^{k+1}(C(\xi^t)) \longrightarrow H^{k+1}(\mathcal{G}_1(\mathbb{R}^m)) \longrightarrow H^{k+1}(\check{X}^t) \longrightarrow \dots$$


We deduce that

$$\text{rank}((\xi^t)^*) = \sum_{k=1}^{+\infty} (-1)^k \left(\dim H^k(\check{X}^t) - \dim H^{k+1}(C(\xi^t)) + \dim H^{k+1}(\mathcal{G}_1(\mathbb{R}^m)) \right)$$

Reminder: $H^1(\mathcal{G}_1(\mathbb{R}^m)) = \mathbb{Z}/2\mathbb{Z}$.

We have to find the image of $(\xi^t)^*: H^1(\check{X}^t) \leftarrow H^1(\mathcal{G}_1(\mathbb{R}^m))$

Let $C(\xi^t)$ be **the mapping cone** of $\xi^t: \check{X}^t \rightarrow \mathcal{G}_1(\mathbb{R}^m)$.



We have a long exact sequence

$$\dots \longrightarrow H^k(\check{X}^t) \longrightarrow H^{k+1}(C(\xi^t)) \longrightarrow H^{k+1}(\mathcal{G}_1(\mathbb{R}^m)) \longrightarrow H^{k+1}(\check{X}^t) \longrightarrow \dots$$

We deduce that

$$\text{rank}((\xi^t)^*) = \sum_{k=1}^{+\infty} (-1)^k \left(\dim H^k(\check{X}^t) - \dim H^{k+1}(C(\xi^t)) + \dim H^{k+1}(\mathcal{G}_1(\mathbb{R}^m)) \right)$$

—————> Can be computed with the persistence algorithm

Conclusion

- We defined persistent Stiefel-Whitney classes,
- Proved stability and consistency results,
- Proposed an algorithm when $d = 1$.

Perspectives:

- Ideas could be extended to other characteristic classes (Euler, Chern, Pontrjagin).
- Need for a triangulation of $\mathcal{G}_d(\mathbb{R}^m)$.

Conclusion

- We defined persistent Stiefel-Whitney classes,
- Proved stability and consistency results,
- Proposed an algorithm when $d = 1$.

Perspectives:

- Ideas could be extended to other characteristic classes (Euler, Chern, Pontrjagin).
- Need for a triangulation of $\mathcal{G}_d(\mathbb{R}^m)$.

Thank you!